

LEONTODON CRISPUS VILL. (COMPOSITAE), A NEW RECORD FROM IRAN

B. Djavadi

Djavadi, B. 1994.12.25: *Leontodon crispus* Vill. (*Compositae*), a new record from Iran. -*Iran. Journ. Bot.* 6 (2): 255-256. Tehran.

Leontodon crispus Vill. subsp. *asper* (Waldst. & Kit.) Rohl. var. *asper*, collected from Marand (NW. of Iran), is reported as a new record from Iran.

Bahereh Djavadi, Department of Botany, Plant Pests and Diseases Research Institute, P.O. Box 1454-19395, Tehran, Iran.

گزارش *Leontodon crispus* Vill. به عنوان گونه جدید برای ایران

باهره جوادی

در بررسی گونه‌های *Leontodon* در هر باریوم ایران، گونه *L. crispus* Vill. که از کوههای میشو DAG (شمال غربی ایران) جمع آوری شده است، برای اولین بار از ایران گزارش می‌شود.



Fig. 1. *Leontodon crispus* subsp. *asper* var. *asper*.

***Leontodon crispus* Vill. subsp. *asper*
(Waldst. & Kit.) Rohl. var. *asper* (Fig.1).**

Azrbaijan: Marand, Payam, Kouhha-ye Mishoudagh (1710-1970 m.), 29 June 1985, Termeh, Moussavi & Tehrani.

This species clearly differs from *L. kotschy* in its 2-4 forked hairs and linear, acute, ± regularly imbricated phyllaries.

The other key characters of this species are:

Leaf indumentum of coarse, stalked, sparingly, branched hairs; outer phyllaries always ciliate-pectinate; dorsal surface of outer phyllaries glabrous or pubescent.

Gen. distri.: SE. Europe. SW. Russia, Crimea, Caucasica and Turkey.