

SEED MORPHOLOGY IMPROVES DATA FROM PHYLOGENETIC RELATIONSHIPS OF MEDICAGOID TRIGONELLA, MEDICAGO AND MELILOTUS (FABACEAE) SPECIES

S. Khandani, M. Assadi, T. Nejdatsari, I. Mehregan & R. Khavarinejad

Received 2017. 05. 20; accepted for publication 2018. 05. 23

Khandani, S., Assadi, M., Nejdatsari, T., Mehregan, I. & Khavarinejad, R. 2018. 06. 30: Seed morphology improves data from phylogenetic relationships of Medicagoid *Trigonella*, *Medicago* and *Melilotus* (Fabaceae) species. -*Iran. J. Bot.* 24 (1): 45-51. Tehran.

Phylogenetic relationship and seed morphological characters of 11 species belong to Medicagoid *Trigonella*, six non Medicagoid and one species of each genera of *Medicago* and *Melilotus* were estimated before based on ITS sequences by using Bayesian inference and maximum parsimony. In this paper, scanning Electron Microscope and stereomicroscope for seed morphological characters were used. Seed morphology results indicated that majority of Medicagoid *Trigonella* species belong to *Bucerates* and *Lunatae* sections placed in two close subclusters and joined together as did by Bayesian analysis results. Both studies confirmed that *Medicago radiata* is sister for these species. Our results confirm that using molecular and morphological characters data are reliable evidence in systematic discriminations of the taxa.

Sima Khandani, Islamic Azad University, Science and Research Branch, Tehran, Iran.-Mostafa Assadi (correspondence < assadi1950@yahoo.com >), Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), P. O. Box 13185-116, Tehran, Iran.-Taher Nejdatsari, Iraj Mehregan & Ramazanali Khavarinejad, Islamic Azad University, Science and Research Branch, Tehran, Iran.

Key words: Fabaceae; Medicagoid; Seed morphology; *Trigonella*; Molecular Phylogeny

ریخت‌شناسی دانه اطلاعات قرابت فیلوژنتیک بین گونه‌های *Melilotus* و *Medicago* و *Medicagoid Trigonella* را با استفاده از مارکرمولکولی ITS تکمیل می‌نماید

سیما خندانی: دانشجوی دکتری سیستماتیک گیاهی، دانشکده علوم دانشگاه آزاد اسلامی، واحد علوم و تحقیقات، تهران
مصطفی اسدی: استاد پژوهش، بخش تحقیقات گیاه‌شناسی، موسسه تحقیقات جنگل‌ها و مراتع کشور، سازمان تحقیقات و آموزش و ترویج کشاورزی، تهران، ایران

ایرج مهرگان: استادیار گروه زیست‌شناسی واحد علوم و تحقیقات دانشگاه آزاد اسلامی، تهران

ظاهر نژادستاری: دانشیار گروه زیست‌شناسی واحد علوم و تحقیقات دانشگاه آزاد اسلامی، تهران

رمضانعلی خاورری نژاد: استاد گروه زیست‌شناسی واحد علوم و تحقیقات دانشگاه آزاد اسلامی، تهران

ارتباط فیلوژنتیکی ۱۱ گونه متعلق به *Medicagoid Trigonella* و ۶ گونه متعلق به *Trigonella* غیر *Medicagoid*، یک گونه از هر کدام از جنس‌های *Medicago* و *Melilotus* با استفاده از توالی منطقه ITS با روش ماکزیمم پارسیمونی و بایسین قبلاً انجام شد. در این مقاله بررسی‌های ریختی دانه توسط میکروسکوپ الکترونی اسکینینگ و استریومیکروسکوپ انجام گرفت. نتایج بدست آمده از ریخت‌شناسی دانه با نتایج حاصل از

مطالعات ITS همخوانی دارد و گونه‌های Medicagoid *Trigonella* در یک دندروگرام با حمایت بالا قرار گرفته و منوفیلیتیک هستند. همچنین نتایج مطالعات مورفولوژی دانه قرار گرفتن گونه‌های بخش‌های *Bucerates* و *Lunatea* متعلق به گروه *Medicagoid Trigonella* در دو زیر خوشه نزدیک به هم را تایید می‌کند. در هردو مطالعه انجام یافته گونه *Medicago radiata* به عنوان گروه خواهری گونه‌های مورد مطالعه در نظر گرفته شد. این نتایج هماهنگی نتایج مورفولوژیکی و مولکولی را در حل اختلافات تاگزونومیکی بیان می‌دارد.

INTRODUCTION

The Fabaceae family is the third largest family of flowering plants that commonly is known as Legum family with about 751 genera and 19000 species that are distributed mainly in temperate and subtropical zones of the world (Mirzaei & al., 2015b). *Trigonella* L. is a large genus with about 135 species from tribe *Trifolieae*. The subtribe *Trigonellinae* from the tribe *Trifolieae* consist of three genera, *Trigonella*, *Medicago* and *Melilotus*. The generic delimitation between *Trigonella* and *Medicago* has long been a problem. Members of this subtribe are united by the consistent presence of pinnately trifoliate leaves and high bootstrap support (99%) in phylogenetic analyses using *matk* (Steele & Wociechoeski, 2003).

Some of the earlier taxonomical studies for delimitation of *Trigonella* and *Medicago* based on floral characters, asymmetric leaves, phenolic variation, stigma morphology, pollen-ovule patterns and seed characters for discriminating between them have been done but with no success (Bena, 2001). Small (1987) based on some floral morphological characters transferred 23 *Trigonella* species including *Bucerates* and *Lunatea* sections to the genus *Medicago* and called them "Medicagoid" species (Gazara & al., 2001). These 23 species have taxonomic problems in the genus *Trigonella* and have been considered as belonging to the genus *Trigonella*. Baum (1968) introduced them as Medicagoid species because of the floral and seed structure similarities among them and *Medicago* species, but based on strong similarities between *Trigonella* and these species in the fruit appearance maintained them in the genus *Trigonella* (Bena, 2001).

Phylogenetic analysis of Medicagoid *Trigonella* L. species based on ITS sequence data showed monophyly of Medicagoid *Trigonella* species while *Bucerates* and *Lunatae* sections were placed in two close subclusters (Khandani & al., 2016b).

Seed Morphological characters in tribe *Trifolieae* have been studied by Lersten & Gunn (1982), Small & al., (1990), Tia (2004), Ceter & al. (2012), Salimpour & al. (2013), Turkie & al. (2013).

The aim of this study is to evaluate and compare the phenetic and phylogenetic relationships within Medicagoid and non Medicagoid *Trigonella* species.

MATERIAL AND METHODS

Taxon Sampling

The materials were taken from dried herbarium specimens and new collection from various parts of Iran during May and June 2014. All of the specimens are deposited in TARI and IAUH. The collected samples were studied and identified by using available floras, monographs and papers. In this study, 19 species including 11 species of Medicagoid *Trigonella* and 6 species of non Medicagoids were selected. Also *Medicago radiata* and *Melilotus officinalis* were chosen as outgroups. (Khandani & al. 2016 a & b).

RESULTS AND DISCUSSION

Seed morphology

Total of 8 quantitative/qualitative characters of seeds including length, width, outline, color, testa texture, homogeneity of cell sizes, testa ornamentation and hilum shape were studied and statistical analysis evaluated. Testa patterns terminology is based on Lersten & Gunn (1982), Small & al., (1990), Pinar & al. (2009), Gunes & Cirpici (2011), Ceter & al. (2012), Gunes (2013), Teixeira & al. (2013), Ozbek & al. (2014) and Mirzaei & al. (2015a) and (Khandani & al. 2016a). *Trifolium orthoceras* and *T. persica* (Medicagoid) have both aculate-tuberculate-verrucate testa ornamentation, circular hilum shape, uniform homogeneity cell size, creased testa texture, oblong outline, but *T. orthoceras* is green color and has length mean 1.76 mm and 0.74 mm width mean. *Trifolium persica* is dark brown color and has mean length 1.95 mm, and mean width 0.69 mm. *Trifolium coerulescens* (non Medicagoid) has reticulate – verrucate testa ornamentation, circular hilum shape, uniform homogeneity cell size, creased testa texture, dark brown color, ovoid outline, length mean 1.47 mm and width mean 0.95 mm. *Trifolium calliceras* (non Medicagoid) has reticulate –verrucate testa ornamentation, elliptic hilum shape, creased testa texture, dark brown color, ovoid outline, length mean 1.77 mm and width mean 1.82 mm. *Trifolium turkemenae* (non Medicagoid) has aculate –reticulate-verrucate testa ornamentation, elliptic hilum shape, uniform homogeneity cell size, creased testa texture, brown color, allantoid –oblong outline, length mean 2 mm and width mean 0.63 mm.

Melilotus officinallis has aculate –tuberculate – verrucate testa ornamentation, elliptic hilum shape, uniform homogeneity cell size, smooth testa texture, dark brown color, ovoid outline, length mean 2.25 mm and width mean 1.52mm.

From Khandani & al. (2016b) can be concluded that phylogenetic tree consists of two main clades (A&B). Clade A includes *Bucerates* and *Lunatae* sections, therefore they are monophyletic taxa. They are also separated from non Medicagoid species and are close to *Medicago radiata* L. than non Medicagoid *Trigonella* species. Other clade includes all non Medicagoid species i. e. *Trigonella coerulescens* (M. B.) Halacsy, *T. elliptica* Boiss., *T. calliceras* Fisch., *T. spruneriana* Boiss., *T. trurkmena* M. Popov and *Melilotus* species.

According to Khandani & al. (2016a), morphological seed analysis observations confirmed the same results based on quantitative and qualitative seed characters (Figs. 1 & 2). Their work introduced phenogram with UPGMA method based on eight traits with highest phenetic correlation ($r > 0.60$) and showed two main clusters. One of them comprises non Medicagoid *Trigonella*, *Medicago radiata* and *Melilotus officinalis* as sister group and the other includes majority of Medicagoid *Trigonella* species forming a separate group. These results are in agreement with previous studies based on seed morphological and molecular data (Tia, 2004; Ceter & al., 2012; Turki & al. 2013; Bena, 2001; Dangi & al. 2016; Downie & al. 1998 and Steele & al. 2010).

In this study, the relation between Medicagoid *Trigonella* species that were shown in both MP and UPGMA analyses (Khandani & al. 2016b) confirmed the works by previous authors (Salimpour & al. 2013). In addition, our work proved close phylogenetic relationship between Medicagoid *Trigonella* species and *Medicago* genus, based on both morphological traits and molecular data. Also, this result is in agreement with some studies on the phylogeny of section *Bucerates*.

Our work shows that both nucleotide sequence data from the internal transcribed spacer (ITS) and seed morphological characters confirm Medicagoid *Trigonella* species separation from other non Medicagoid *Trigonella* species, which was suggested by Small (1987), based on floral characters. These results agree with the treatment in Flora Iranica (Rechinger 1984) which separated majority of Medicagoid *Trigonella* species in *Bucerates* section, such as two close subclusters based on seed morphology (Khandani & al. 2016a) and one cluster based on ITS sequences data (Khandani & al. 2016 b).

ACKNOWLEDGEMENT

We would like to thank the authorities of Islamic Azad University Science and Research Branch, Tehran, Iran and Research Institute of Forest and Rangeland, Iran for providing the facilities necessary to carry out the study.




Fig. 1. SEM microphotographs of *Trigonella* seeds: A-B, *Trigonella coerlescens*; C-D, *T. orthoceras*; E-F *T. persica*.




Fig. 2. SEM microphotographs of *Trigonella* and *Melilotus*. A-B, *Trigonella turkemena*; C-D, *T. calliceras*; E-F, *Melilotus officinalis*.




Fig. 3. Bayesian consensus phylogram tree. Numbers above branches are Bayesian posterior probabilities. Numbers below branches are bootstrap values) and SEM microphotography of *Trigonella* and *Melilotus*.

REFERENCES

- Bena, G. 2001: Molecular phylogeny supports the morphologically based taxonomic transfer of the "Medicagoid" *Trigonella* species to the genus *Medicago* L.-Plant Syst. Evol. 229: 217-236
- Ceter, T., Pinar, M., Akan, H., Ekici, M. & Aytac, Z. 2012: Comparative seed morphology of *Trigonella* L. species (Leguminosae) in Turkey. -Afr. J. Agric. Res. 7 (3): 509-522.
- Dangi, R., Tamhankar, S., Choudhary, R.K. & Rao, S. 2016: Molecular phylogenetics and systematics of *Trigonella* L. (Fabaceae) based on nuclear ribosomal ITS and chloroplast *trnL* intron sequences.-Genetic. Resour. Crop. Evol. 63(1):79-96.
- Downie, S.R., Katz-Downie, D.S., Rogers, E.J., Zvjewski, H. L. & Small, E. 1998: Multiple Independent losses of the plastid *rPoC1* intron in *Medicago* (Fabaceae) as inferred from phylogenetic analyses of nuclear ribosomal DNA internal transcribed spacer sequences. -Can. J. Bot. 76 (5): 791-803.
- Gazara, M., Kamel, W., Haider, A. 2001: Cladistic analysis of the genera: *Trifolium*, *Trigonella* and *Melilotus* (Fabaceae; Papilionaceae). -Egypt. J. Biol. 3: 161-170.
- Gunes, F. & Cirpici, A. 2011: Seed characteristics and testa textures some taxa of genus *Lathyrus* L. (Fabaceae) from Turkey. -Int. J. Agric. Biol. 13 (6): 1814-9596.
- Gunes, F. 2013: Seed characteristics and textures of *Pratensis*, *Orobon*, *Lathyrus*, *Orobastrum* and *Cicercula* sections from *Lathyrus* (Fabaceae) in Turkey. -Plant Syst. Evol. 299: 1935-1953.
- Khandani, S., Assadi, M., Nejdassattari, T. & Mehregan I. 2016a: Phenetic analysis of genera Medicagoid *Trigonella*, *Medicago* and *Melilotus* (Fabaceae) on seed coat in Iran.-Biodiversitas vol 17, No. 1: 162-171.
- Khandani S., Assadi M., Mehregan I. & Nejdassattari T. 2016 b: Phylogenetic analysis of Medicagoid *Trigonella* L. species based on ITS sequence data. Ambient Science. vol. 3: 35-39.
- Lersten, NR., Gunn, CR. 1982: Testa characters in tribe *Vicieae* with notes about tribes *Abreae*, *Cicereae*, and *Trifolieae* (Fabaceae).-U. S. Department of Agriculture Technical Bulletin. No. 1667: 1-40.
- Mirzaei, L., Assadi, M., Nejdassattari, T. & Mehregan, I. 2015a: Seed surface morphology in some species of the *Colutea* L. (Fabaceae) in Iran. -Adv. Biores. 6 (5): 105-109.
- Mirzaei, L., Assadi, M., Nejdassattari, T. & Mehregan, I. 2015b: Comparative seed and leaf micromorphology of *Colutea* species (Fabaceae) from Iran. -Environ. Exp. Biol. 13: 183-187.
- Ozbek, F., Ozbek, MO. & Ekici, M. 2014: Morphological, anatomical, pollen and seed morphological properties of *Melilotus bicolor* Boiss & Balansa (Fabaceae) endemic to Turkey. - Aust. J. Crop. Sci. 8 (4): 543-549.
- Pinar, MN., Duran, A., Ceter, T., Tug, N. 2009: Pollen and seed Morphology of the Genus *Hesperis* L. (Brassicaceae) in Turkey. -Turk. J. Bot. 83-96.
- Rechinger, K.H. 1984: *Trigonella*. -Flora Iranica 157: 207-253. Graz.
- Salimpour, F., Safiedin, M., Sharifnia, F. 2013: Phylogenetic relationships between *Trigonella* species (*Bucerates* section) using ITS markers and morphological traits. -J. Bio. Env. Sci. 3 (12): 116-124.
- Small, E., Brookes, B., Lassen, P. 1990: Circumscription of the genus *Medicago* (Leguminosae) by seed characters. -Can. J. Bot. 68: 613-629.
- Small, E. 1987: Generic changes in *Trifolieae* subtribe *Trigonellinae*. -Advances in Legume. Vol. 3. - Royal Botanic Gardens, Kew. 169-181.
- Steele, KP. & Wojciechowski, MF. 2003: Phylogenetic analyses of tribes *Trifolieae* and *Vicieae*, based on sequences of the plastid gene *matk* (Papilionoideae: Leguminosae). -Advances in Legume Systematics, part 10, Higher Level Systematics. Royal Botanic Gardens, Kew.
- Steele, KP., Ickert-Bond, M., Zarre, S., Wojciechowski, MF. 2010: Phylogeny and character evolution in *Medicago* (Leguminosae). Evidence from analyses of plastid *trn/matk* and nuclear GA3OX1 sequences. Am. J. Bot. 97: 1142-1155.
- Teixeira, de QR., Azevedo, de TAM., Lewis, GP. 2013: Seed morphology: an addition to the taxonomy of *Tephrosia* (Leguminosae, Papilionoideae, Millettieae) from South America. Plant Syst. Evol. 299: 459-470
- Tia, WK. 2004: Tribe *Trifolieae*: Evidence from seed characters. Pak. J. Biol. Sci. 7 (7): 1287-1302.
- Turki, Z., El-Shayeb, F., Abozeid, A. 2013: Seed morphology of some *Trigonella* L. species (Fabaceae) and its taxonomic significance. Int. J. Sci. Res. 2319-7064.