

CHROMOSOME NUMBERS FOR *RANUNCULUS DOLOSUS* AND *SILENE CONOIDEA* FROM IRAN

F.S. Ghasemi, A. Jalili, S. S. Mirzadeh Vaghefi & S. Ashrafi

Received 2015. 02.15; accepted for publication 2015. 06. 06

Ghasemi, F.S., Jalili, A., Mirzadeh Vaghefi, S. S. & Ashrafi, S. 2015. 06. 30: Chromosome numbers for *Ranunculus dolosus* and *Silene conoidea* from Iran.- *Iran. J. Bot.* 21 (1): 61-63. Tehran.

Chromosome numbers are reported for *Ranunculus dolosus* (Ranunculaceae), $2n=32$ and *Silene conoidea* (Caryophyllaceae), $2n=24$. These species were collected from NW Iran in W Azarbaijan Province. Idiograms were prepared for both species. Symmetrical karyotypes are placed in class 1B in *Ranunculus dolosus* and class 1A in *Silene conoidea*. Karyotype formula are $8m+6sm+2st$ in *Ranunculus dolosus* and $12m$ in *Silene conoidea*.

Farzaneh sadat Ghasemi (correspondence <f. ghsemi@rifr-ac.ir>), Adel Jalili, Saeede sadat Mirzadeh Vaghefi & Soheila Ashrafi, Research Institute of Forests & Rangelands, P.O.Box 13185-116, Tehran, Iran.

Key words: Karyotype; chromosome number; *Silene conoidea*; *Ranunculus dolosus*; Iran

اعداد کروموزومی گونه‌های *Ranunculus dolosus* و *Silene conoidea* از ایران

فرزانه سادات قاسمی، کارشناس ارشد موسسه تحقیقات جنگل‌ها و مراتع کشور، تهران، ایران.

عادل جلیلی، استاد پژوهش موسسه تحقیقات جنگل‌ها و مراتع کشور، تهران، ایران.

سعیده سادات میرزاده واقفی، کارشناس ارشد موسسه تحقیقات جنگل‌ها و مراتع کشور، تهران، ایران.

سهیلا اشرفی، کارشناس ارشد موسسه تحقیقات جنگل‌ها و مراتع کشور، تهران، ایران.

عدد کروموزومی و کاریوتیپ گونه‌های: *Ranunculus dolosus* (Ranunculaceae) ($2n=32$) و *Silene conoidea* (Caryophyllaceae) ($2n=24$) از ایران گزارش می‌شوند. رویشگاه‌های طبیعی این گونه‌ها شمال غرب ایران است. ایدیوگرام گونه‌ها ارائه شدند. تقارن کاریوتیپی در گونه *Ranunculus dolosus* 1B و در گونه *Silene conoidea* 1A می‌باشند. فرمول کاریوتیپی در گونه *Ranunculus dolosus* $8m+6sm+2st$ و در گونه *Silene conoidea* $12m$ می‌باشند.

INTRODUCTION

Cytological data are of great importance for understanding of relationship and evolution in angiosperms (Ehrendorfer & al. 1968; Keshavarzi 2012). Determination of chromosome numbers and karyotypes of the native Iranian plants are the major aim in this study. Chromosome numbers of some Iranian plants have been reported by Ghahramaninejad & al (2013). In this work the cytological data for *Ranunculus dolosus* (Ranunculaceae), an endemic species and *Silene conoidea* (Caryophyllaceae) are provided. These species were collected from wetland habitats in NW of Iran. *Ranunculus dolosus* is an endemic of Iran.

MATERIALS AND METHODS

For cytological study, rootlets were collected from

germinated seeds on wet filter paper in petri dishes at room temperature, when they reached 1–1.5 cm in length, rootlets were collected. The material was pretreated in alphabromonaphtalene for 2 hour, fixed in solution of ethanol- acetic acid (3: 1) for 4 hours. The meristems stained in hematoxylin and squashed in a drop of 45% acetic acid solution. Metaphase plates were examined with light microscope, Olympus BH2-RFCA under oil immersion (x100).

RESULTS

Ranunculaceae

Ranunculus dolosus Fisch. & C. A. Mey

Specimen studied: East Azarbaijan: Boostan Abad, After Said Abad Ghoorigol, 1869 m, 30.07.2013, N $37^{\circ}.54'.02.40''$, E $46^{\circ}.42'.5.11''$, Ranjbari, 8234 East Azarbaijan Herbarium.

The genus *Ranunculus* have about 50 annual and perennial herbaceous species in Iran (Pakravan 2012). Basic number of chromosomes in *Ranunculus* species are $x=7$ and $x=8$ (Ovidio & Marchi 1990). The somatic chromosome number $2n=32$ is reported here for the first time for *Ranunculus dolosus* (fig. 1). Idiogram of *Ranunculus dolosu* is presented (fig 2). Symmetrical karyotype were in class 1B (Stebbins 1971). Karyotype formula was $8m+6sm+2st$.

Caryophyllaceae

Silene conoidea L.

Specimen studied: West Azarbaijan, Oroomieh, Salmas between village Koozerash and Haji Jaghan, 2281 m, 20.07.2013, N $38^{\circ}.08'.40.9''$, E $44^{\circ}.27'.25.3''$, Ashrafi 101602 TARI.

The genus *Silene* has about 100 species in Iran (Nejati Edalatiyan & al 2011) which are herbaceous annuals and perennials. Chromosome number of *S. conoidea* was previously reported as $2n=20, 24, 30, 48, 72$ and $n=20$ (Abdel Bari 1973, Dong 2011, Jeelani 2011). Also, chromosome number $2n=24$ has been reported in a previous work (Flora of North America Editorial Committee 2005). In Iran the previous report on *S. conoidea* collected from Chaharmahal va Bakhtiari Province shows $2n=20$ (Aryavand & Favarger 1980) The somatic chromosome number $2n=24$ is reported here for this species (fig. 1). Chromosomes are moderately symmetrical, in class 1A (Stebbins 1971). Idiogram of *Silene conoidea* is presented. Karyotype formula is $12m$, (fig. 2).

REFERENCES

- Abdel Bari, E. 1973: Cytological studies in the genus *Silene* L.- New phytologist 72:833-838
- Aryavand, A.& Favarger, C. 1980: Contribution à l'étude Cytotaxonomique des Caryophyllaceae de l'Iran. -Rev. Biol. Ecol. Méditerranéenne t. VII, no 1: 15-26.
- Dong, L., Dan, L., Bo, X., Ze-Long, N, Hang, S & Zhi-Min, L. 2011: A karyological study of six species of *Silene* L. (Caryophyllaceae) from the Hengduan Mountains, SW China, Caryologia 64 1:10-13.
- Ehrendorfer, F., Krendl, F., Habeler, E. and Sauer W. 1968: Chromosome Numbers and Evolution in Primitive Angiosperms.- Taxon, vol. 17, No. 4: 337-353
- Flora of North America Editorial Committee (Ed.). 2005: Flora of North America: *Magnoliophyta*: Caryophyllidae, pt. 2 (Vol. 5). -Oxford University Press.
- Ghahramaninejad, F., Najad Falatoury, A., Mahmoodi, M., Fereidounfar, S & Hoseini, E., 2013: Plant chromosome book of Iran, Biology home press, 194pp.-Tehran.
- Jeelani, S., Rani, S., Kumar, S. & Gupta, R., 2011: Meiotic studies in some members of Caryophyllaceae Juss. from the Western Himalayas.-Acta Biologica 53;1:86-95
- Keshavarzi, M. 2012: Chromosome numbers for some *Eremopyrum* Jaub.& Spach(Poaceae) species of Iran.- Iran. J. Bot.18 (1):94-97.Tehran.
- Nejati Edalatiyan., M., Joharchi, M. & Ghahramaninejad, F., 2011: *Silene ferdowsii* (Caryophyllaceae) a new species from Iran.- Ann. Bot. Fennici 48:155-158.
- Ovidio, R and Marchi, P., 1990: DNA Content, Karyotype Structure Analysis and Karyotype Symmetry in *Ranunculus* L.(Ranunculaceae). Italian Species Belonging to Sections *Flammula* (Webb) Benson and *Micranthus* (Ovcz.) Nyarady.- Caryologia Vol. 43, no. 2: 99-115.
- Pakravan, M. 2012: A new species of the genus *Ranunculus* from Iran.- Science Asia 38: 419-421.
- Stebbins, G., L. 1971: Chromosome Evolution in Higher Plants.- Edvard Arnold Publisher, London.
- Yang, Q.E. 2001: Cytology of eleven species in the genus *Ranunculus* L. and five in its four related genera from China.- Acta Phytotaxonomica Sinica, 39(5):405-422.




Fig.1. Somatic metaphases in *Ranunculus dolosus* (left); *Silene conoidea* (right).




Fig 2. Idiograms of *Ranunculus dolosus* (above) *Silene conoidea* (below).