

SAMERARIA STENOPHYLLA (BRASSICACEAE), A NEW RECORD FOR THE FLORA OF IRAN

H. Moazzeni, S. Sajedi & M. R. Joharchi

Moazeni, H., Sajedi, S. & Joharchi, M.R. 2006 08 01. *Sameraria stenophylla* (Brassicaceae), a new record for the flora of Iran. – *Iran. J. Bot.* 12(1): 63-66. Tehran.

Sameraria stenophylla is reported as a new record from E Iran. This species is easily distinguished from its closest relative, *Sameraria elegans*, by having larger petals and stem leaves which are sagittate-cordate at base. A geographical distribution map and an illustration of the species are given.

Hamid Moazzeni, School of Biology, College of Science, Tehran University, Tehran, Iran, P. O. Box: 14155-6455, e-mail: moazzeni@khayam.ut.ac.ir – Sepideh Sajedi, Botany Department, Plant Pests and Diseases Research Institute, P.O.Box 1454.Tehran 19395. Iran. – Mohammad Reza Joharchi, Ferdowsi University of Mashhad Herbarium (FUMH), Mashhad, Iran.

Key words. *Sameraria*, Brassicaceae, new record, Iran.

گزارش *Sameraria stenophylla* به عنوان گونه جدیدی برای ایران

حمید موذنی، سپیده ساجدی و محمد رضا جوهرچی

گیاه *Sameraria stenophylla* به عنوان گونه جدیدی از شرق ایران گزارش می شود. این گونه به سادگی با داشتن گلبرگ‌های بلند و سرنیزه‌ای بودن قاعده برگ‌های ساقه‌ای از گونه مجاور خود یعنی *Sameraria elegans* قابل تفکیک است. نقشه پراکنش این گونه و نزدیکترین خویشاوند آن به همراه تصویری از این گونه ارائه می‌شود.

Introduction

Iran, with about 120 genera representing 358 species of Brassicaceae (Akhani 2003), is one of the most important diversification centers of the family in the world. After the publication of the Brassicaceae for Flora Iranica (Hedge 1968), several new species or new combinations have been proposed for Iran (see for example: Sajedi & al. 2004, Akhani 2003, Khosravi 2003, Kavousi 2001, Sonboli 2000, Moussavi 1985, Mozaffarian 1985, Assadi 1983). *Sameraria* Desv. is a genus of about nine species distributed solely in Central and SW Asia (Appel & Al-Shehbaz 2003). It is closely related to *Isatis* and shows several characters in common with it, namely the habit, leaves, inflorescence and flowers. However, it is separated from *Isatis* by having silique and/or silicule fruits possessing short and distinct style, instead of mostly silique fruits and sessile stigmas.


In the framework of preparing a revision of *Sameraria* and its allies as the M.Sc. thesis by the first author, we could identify *Sameraria stenophylla* among the material deposited in the Iranian herbaria, including Ferdowsi (Mashhad) University (FUMH). This species has not previously been reported from Iran. This new record elevates the number of *Sameraria* species distributed in Iran to five out of the six species reported by Hedge (1968) for entire Flora Iranica area.

Sameraria stenophylla Rech. f. & Koeie (sect. *Sameraria*), Fig. 1.

Specimens seen:

IRAN: Prov. Semnan, 6 km W Eyvanaky, 1 May 1974, Iranshahr & Riedl s.n. (IRAN). Prov. Khorassan: Shahrud to Sabzevar, Delbar village, 1100 m, 28 April 1975, Iranshahr 30756 (IRAN). Khorassan, E. Ghain, Petergan, Yazdan village, 450 m, Joharchi & Zangooei 16005 (FUMH). **AFGHANISTAN.** Prov. Kandhar, Akhundzadeh, 19 km SW Shar-e Safa on the road to Kandahar, 1170 m, 15 April 1972, Anders 8374 (MSB).

Annuals 15–40 cm tall, glabrous except for fruits, branched at base. Basal leaves obovate to oblong, 2–8 × 1–1.5 (–2.2) cm, lobed to entire at margins, obtuse to rounded at tip, cuneate at base; petiole 0–1 cm long. Cauline leaves linear to oblong, sagittate or cordate at base, narrowing towards apex. Fruiting pedicel 7–10 mm long, thin, deflexed. Sepals oblong, 3.8–4.5 × 1.5 mm, glabrous, violet to pale yellow. Petals 6–7 × 2–2.5 mm, obovate, cuneate at base. Long filaments 3.5–3.8 mm long, short filaments ca. 2.5 mm long. Fruit indehiscent, 1-seeded, ca. 15 × 15 mm, orbicular to broadly ovate, broadly winged at margins, suture wings absent, below the locule cylindrical, emarginate to cordate at base, densely villous; style 1–3 mm long.



Map 1. Distribution of *Sameraria stenophylla* (■) and *S. elegans* (●) in Iran.

General distribution (map. 1):

According to Hedge (1968), *S. stenophylla* is endemic to SW Afghanistan, the area around Jija in Prov. Farah. We found also a second material of the same species in the herbarium of Institute of Systematic Botany in Munich collected by O. Anders in Afghanistan (deposited in MSB) from Prov. Kandhar, near Shar-e Safa. So, the species is distributed much wider in Afghanistan than previously known, and its presence at eastern borders of Iran is not surprising.

This species is closely related to *Sameraria elegans* Boiss. The differences between these species are summarized in table 1.

Table 1. Diagnostic characters useful in separating *Sameraria stenophylla* and *S. elegans*.

Taxon→	<i>S. stenophylla</i>	<i>S. elegans</i>
Character↓		
Petals	6–7 × ca. 2 mm	ca. 4 × 1 mm
Style	1–3 mm	0.5–1.5 mm
Cauline leaves at apex	attenuate	obtuse
Cauline leaves at base	sagittate-cordate	rounded-auriculate

Basal leaves	lobed to entire	entire
--------------	-----------------	--------

ACKNOWLEDGMENT

Authors are sincerely grateful to Mr. M. Mehranfard for drawing of the illustration and to Ihsan A. Al-Shehbaz for verification of the identity and for critical review of the manuscript. We are also grateful to Dr. Shahin Zarre for his useful comments on the manuscript.

References

- Akhani, H. 2003: Notes on the flora of Iran, IV, two new records and synopsis of the new data on Iranian Cruciferae since Flora Iranica. -Candollea 58(2): 369–385.
- Appel, O. & Al-Shehbaz, I. A. 2003: Cruciferae. In K. Kubitzki (editor), Families and Genera of Vascular Plants 5: 75–174. -Springer-Verlag, Berlin and Heidelberg.
- Assadi, M. 1983: Plants from N. Azarbaijan, new to Iran. -Iran. J. Bot. 2 (1): 1–8.
- Hedge, I. C. 1968. *Sameraria* in Rechinger K.H. (ed.) Flora Iranica 57: 93–96. – Graz, Austria: Akademische Druck- und Verlagsanstalt.

- Jafri, S.M.H. 1973: Brassicaceae in Nasir E. & Ali S. I. (eds.) Flora of West Pakistan 55: 1–308. -Karachi: Pakistan.
- Kavousi, K. 2001: Notes on the plant family Cruciferae in Iran, new taxa and new records. -Iran. J. Bot. 9: 47–54.
- Khosravi, A. R. 2003: A reconsideration of *Hesperis leuoclada* Boiss. (Cruciferae). -Iran. J. Bot. 10(1): 15–23.
- Mousavi, M. 1985: *Fibigia thesigeri* (Cruciferae), a new record from Iran. -Iran. J. Bot. 3: 63–64.
- Mozaffarian, M. 1985: New species and new plant records from Iran. -Iran. J. Bot. 3: 81–86.
- Rechinger, K. H. 1968: Cruciferae in Rechinger, K. H. (ed.) Flora Iranica 57: 1–372. – Graz, Austria: Akademische Druck- u. Verlagsanstalt.
- Sajedi, S., Sharifnia, F. & Sonboli, A. 2004: Verification of *Isatis tinctoria* in Iran.-Rostaniha 5: 51–52.
- Sonboli, A., Assadi, M., Zehzad, B. 2000: *Sterigmostemum laevicaule* Bornm., a synonym of *Erysimum crassicaule* (Boiss) Boiss. (Cruciferae). - Willdenowia 30: 359–360.