

A NEW ISOLATED SPECIES OF ASTRAGALUS SECT. CAPRINI (FABACEAE) FROM KHORASSAN

Maassoumi, A. A., Safavi, H. & Nasseh, Y.

Maassoumi, A. A., Safavi, H. & Nasseh, Y. 2001 12 15: A new isolated species of *Astragalus* sect. *Caprini* (*Fabaceae*) from Khorassan. -*Iran. Journ. Bot.* 9 (1): 63-64. Tehran.

A. new isolated species of *Astragalus*, sect. *Caprini* (*Fabaceae*) from Khorassan, *A. zangooianus* is described as new from E. Iran, Khorassan province.

Ali Asghar Maassoumi, Research Institute of Forests and Rangelands, P. O. Box 13185-116, Tehran, Iran. -Hekmat Safavi & Yasaman Nasseh, University of Ferdowssi Herbarium.

Key words. *Astragalus*, *Caprini*, Iran, new species, Khorassan.

یک گونه جدید منفک گون از بخش **Caprini** از خراسان
علی اصغر معصومی، حکمت صفوی و یاسمون ناصح

در میان گونه‌های شناخته شده فراوان متعلق به بخش *Caprini* از استان خراسان، گونه
A. zangooianus به عنوان یک گونه جدید منفک از شرق ایران، استان خراسان شرح داده
می‌شود.

Introduction

In the course of identification of unnamed material, for preparation of the second version of the genus *Astragalus* in the form of Flora of Iran, *Astragalus zangooianus* was recognized as a distinct isolated new species from sect. *Caprini*.

Astragalus zangooianus Maassoumi, Safavi & Nassee, sp. nov. e sect. *Caprini*.

Similar to *A. remotijugus* Boiss. & Hohen., but it differs in having the violet calyx (not yellow); standard deeply violet (not yellow), limb of keel with minute teeth, pods at the base with a long stipe ca. 5 mm long (not sessile).

Planta perennis, acaulis, ca. 15-25 cm alta, glaberrima. Stipulae ovato-lanceolatae, ca. 15 mm longae, glabrae, basi ca. 5 mm pectiolo adnatae, inter se liberae. Folia 15-25 cm longa, pectiolo ca. 4-8 cm longo; foliola 6-8 juga, remota, obovata ad orbicularia, apice retusa vel emarginata, raro obtusa. Inflorescentiae pedunculatae, pedunculis ca. 6-7 cm longis, 3-4 (-6) florae. Calyx tubulosus, purpureus, ca. 15-17 mm longus, basi distinete gibbosus, dentibus linearo-

lancolatis, ca. 5 mm longis. Corolla intense violacea. Vexillum ca. 20-22 mm longum et 10 mm latum; lamina oblongo-elliptica, antice rotundata. Alae ca. 20 mm longae; laminae ellipticae, apicem versus leviter dilatatae, basi auriculatae, auriculae ca. 4 mm longae. Carina ca. 16 mm longa; lamina curvato-triangularis, in margine superiore denticulis minutis provisa. Ovarium glabrum, basi stipite 6-7 mm longo suffultum, stylo sub stigmate glabro. Legumen basi stipite ca. 8 mm longo suffultum, cum stipe ca. 4 cm longum, ca. 1 cm latum, dorso-ventro applanatum, biloculare.

Holotypus. Khorassan, Kashmar to Neyshabur, ca. 55 km from Kashmar, 1850 m, 11. 5. 1991, Joharchi & Zangoei 20083 (TARI; Iso.: Herbarium of Ferdousi University, Mashhad). —*Paratypus.* Khorassan, Torbat-e Heidariyeh, Kuhay-e Khargh, 1750-2200 m, 2.4.1994, Paryab & Mossavi 3050 (Research Centre of Khorassan).

Species named in the honour of Mr. Zangoei, collector of the University Research Center.