

# THE STUDY OF HETEROCYSTOUS CYANOBACTERIA FROM PADDY FIELDS IN KALAT NADERI DISTRICT IN NORTH-EAST OF IRAN

E. Aslani, H. Riahi, Z. Shariatmadari & F. Bazzi

Received 2013. 12. 25; accepted for publication 2014. 02. 25

Aslani, E., Riahi, H., Shariatmadari, Z. & Bazzi, F. 2014. 12. 31: The Study of Heterocystous Cyanobacteria from Paddy Fields in Kalat Naderi Region in North-East of Iran.- *Iran. J. Bot.* 20 (2): 257-264. Tehran.

Heterocystous cyanobacteria represent an interesting group of photosynthetic prokaryotes which are capable of N<sub>2</sub> fixation and are present abundantly in rice fields. The aim of this study was the identification and purification of cyanobacteria from rice fields in the Kalate Naderi, which is located in north-east of Khorasan Razavi Province. To approach this aim, four paddy fields were selected to investigate heterocystous blue-green algae. In total, 30 taxa were identified. These species belong to genera such as *Nostoc*, *Anabaena*, *Cylindrospermum* and *Calothrix*. Among these genera, *Nostoc* was found to be dominant genus in all samples. Six species, including *Anabaena fuellbornii* Schmidle, *Anabaena inaequalis* Bornet & Flahault, *Nostoc pruniforme* C. Agardh ex Bornet & Flahault, *Nostoc amplissimum* Setchell, *Nostoc piscinale* Kützing ex Bornet & Flahault, and *Calothrix fusca* (Kützing) Bornet & Flahault were identified as new records for Iran.

Elahe Aslani, Hossein Riahi (correspondence < h-riahi@cc.sbu.ac.ir >), Zeinab Shariatmadari & Fateme Bazzi, Faculty of Biosciences, University of Shahid Beheshti, Evin, Tehtan, Iran.

**Key words:** Algal flora; Heterocystus cyanobacteria; Kalat Naderi; Paddy field; Khorassan; Iran

مطالعه سیانوباکتری‌های دارای هتروسیتست مزارع برنج کلات نادری واقع در شمال شرق ایران

الهه اصلانی، دانشکده علوم زیستی، دانشگاه شهید بهشتی، تهران

حسین ریاحی، استاد دانشکده علوم زیستی، دانشگاه شهید بهشتی، تهران

زینب شریعتمداری، استادیار دانشکده علوم زیستی، دانشگاه شهید بهشتی، تهران

فاطمه بازی، دانشکده علوم زیستی، دانشگاه شهید بهشتی، تهران

سیانوباکتری‌های تولیدکننده هتروسیت گروه ویژه‌ای از پروکاریوت‌های فتوسنتزکننده هستند که قادرند نیتروژن اتمسفری را تثبیت کنند و در مزارع برنج به فراوانی یافت می‌شوند. هدف از این مطالعه خالص‌سازی و شناسایی سیانوباکتری‌های مزارع برنج ناحیه کلات نادری از استان خراسان رضوی است. در راستای نیل به این هدف، فلور جلبکی ۴ ایستگاه واقع در این ناحیه مورد بررسی قرار گرفت و در مجموع ۳۰ تاکسون شناسایی شد. گونه‌های شناسایی شده مربوط به جنس‌هایی از قبیل *Nostoc*، *Anabaena*، *Cylindrospermum* و *Calothrix* بودند. در میان این جنس‌ها، جنس *Nostoc* به عنوان نمونه غالب موجود در تمام ایستگاه‌ها شناسایی شد. همچنین شش گونه ذیل به عنوان گونه‌های جدید برای ایران گزارش می‌گردند:

*Anabaena fuellbornii* Schmidle, *Anabaena inaequalis* Bornet & Flahault, *Nostoc piscinale* Kützing ex Bornet, *Nostoc pruniforme* C. Agardh ex Bornet & Flahault, *Nostoc amplissimum* Setchell, *Calothrix fusca* (Kützing) Bornet & Flahault.

## INTRODUCTION

Cyanobacteria are oxygenic photosynthetic microorganisms which are widely distributed over a diverse range of habitats (Geitler 1932; Frey 1933;

Desikachary 1959). Among this group of algae, several taxa with positive and negative environmental impacts have been identified. For example, a significant number of Nostocaceae family are capable of nitrogen fixation,




Fig. 1. Sampling sites in Kalat Naderi.

while some of them are known as harmful algae by producing toxins in aquatic ecosystems (Galhano & al. 2011). Nitrogen fixing cyanobacteria are the most important group of these algae with positive environmental effect. They are found in both free living as well as symbiotic forms (Ezhilarasi & Anand 2009). In addition to the nitrogen fixation ability, they appeared to be a rich source of many useful compounds and play an important role in improving soil condition. Consequently, cyanobacteria can increase plant growth and yield as a natural biofertilizer (Song & al. 2005).

The order Nostocales is one of the most important groups of diazotrophic cyanobacteria. This order includes filamentous cyanobacteria that are capable of cell differentiation in heterocysts, akinetes and vegetative cells (Castenholz 1989; Anagnostidis & Komárek 1989). Using morphological characters is considered important way to identify heterocystous cyanobacteria. These include characters such as morphology of filaments, especially different shapes of vegetative cells, heterocysts and akinetes. Up to now, several species from different genera of Nostocacean cyanobacteria were reported from terrestrial habitats of Iran (Nowruzi & Ahmadimoghdam 2006; Saadatnia & Riahi 2009; Siahbalaie & al. 2011; Shariatmadari & al. 2013). Nevertheless, there are limited studies on the cyanobacterial flora of paddy soils of northeast of Iran. The Kalat Naderi district is located in northeast of Iran in Khorasan Razavi Province. The aim of the present work was to study Nostocacean cyanobacteria from rice fields of Kalat Naderi district.

## MATERIALS AND METHODS

This study is focused on Kalat Naderi district of Khorasan Razavi Province. The Kalat Naderi township is located in the north-east of Iran and between 59° 9'E

to 60° 27' E longitude and 36° 24' N to 37° 17' N latitude (fig.1). Soil samples were collected from four sites (Amirabad, Sarrood, Qaleno, Kalat) in April 2011. The depth of sampling was 5 cm (Stanier & al. 1971). The collected soil samples were transferred to sterilized nitrate free BG-11 medium. The Petri dishes were placed in a culture chamber (Noorsanat Tissue Culture Shelf, Model SB5520) at 25±5°C and a 12/12h light-dark cycle under fluorescent illumination of 2000 to 2500 Lux for two weeks. After colonization, the cyanobacterial strains were studied by optical microscopy (Olympus, Model BM-2) and different taxa were identified based on morphological characteristics using standard key books and articles (Desikachary 1959; Prescott 1970; Komárek 2005). Morphology of filaments, shape and dimensions of vegetative cells, heterocysts and akinetes were some of the characters used for identification of these taxa.

## RESULTS

In the present study, 30 taxa of heterocystous cyanobacteria were identified and recorded from Kalat Naderi district (figs. 2, 3, 4). Twenty six species belonging to three genera of Nostocaceae and one genus with four species of Rivulariaceae were listed in table 1. The genus *Nostoc* with 15 species were predominant and comprising 50% of all recorded taxa (table 2). Six species of identified taxa, *Anabaena fuellebornii*, *A. inaequalis*, *Nostoc amplissimum*, *N. pruniforme*, *N. picinale* and *Calothrix fusca* were considered as new records for Iran (fig. 2).

### Description of the new records Nostocaceae

1. *Anabaena inaequalis* (Kütz.) Bornet & Flahault 1888 (fig. 2A)

Table 1. List of cyanobacterial species recorded from paddy fields of Kalat Naderi. \* indicates new records for Iran.

Taxa	Sites			
	1	2	3	4
<i>Anabaena vaginicola</i> F.E.Fritsch & Rich	-	+	+	+
<i>Anabaena torulosa</i> Lagerheim ex Bornet & Flahault	-	+	+	+
<i>Anabaena oscillarioides</i> Bornet & Flahault	-	+	+	+
<i>Anabaena orientalis</i> S.C.Dixit	-	-	+	-
<i>Anabaena portoricensis</i> Gardner	-	-	+	-
<i>Anabaena iyengarii</i> Bharadwaja	-	-	+	-
* <i>Anabaena fuelebornii</i> Schmidle	-	-	+	-
* <i>Anabaena inaequalis</i> Bornet & Flahault	-	-	-	+
* <i>Nostoc piscinale</i> Kützing ex Bornet & Flahault	-	-	+	-
* <i>Nostoc entophyllum</i> Bornet & Flahault	+	-	-	-
<i>Nostoc amplissimum</i> Setchell	-	-	+	-
<i>Nostoc alatosporum</i> Sant'Anna	-	-	-	-
<i>Nostoc muscorum</i> C.Agardh ex Bornet & Flahault	+	-	+	-
<i>Nostoc rivulare</i> Kützing ex Bornet & Flahault	-	-	+	-
<i>Nostoc commune</i> Vaucher ex Bornet & Flahault	-	+	-	-
* <i>Nostoc pruniforme</i> C.Agardh ex Bornet & Flahault	-	-	+	-
<i>Nostoc sphaericum</i> Vaucher ex Bornet & Flahault	-	-	+	-
<i>Nostoc spongiaeforme</i> C.Agardh ex Bornet & Flahault	-	-	+	+
<i>Nostoc linckia</i> (Roth) Bornet ex Bornet & Flahault	-	-	+	-
<i>Nostoc ellipsosporum</i> Rabenhorst ex Bornet & Flahault	-	-	+	+
<i>Nostoc punctiforme</i> (Kützing) Horjot	-	-	+	-
<i>Nostoc calcicola</i> Brébisson ex Bornet & Flahault	+	+	+	+
<i>Nostoc verrucosum</i> Vaucher ex Bornet & Flahault	+	-	-	-
<i>Cylindrospermum sphaericum</i> B.N.Prasad	-	-	+	-
<i>Cylindrospermum stagnale</i> Bornet & Flahault	-	-	+	-
<i>Cylindrospermum muscicola</i> Kützing ex Bornet & Flahault	-	-	+	-
<i>Calothrix elenkinii</i> Kossinskaja	-	+	+	+
<i>Calothrix dolichomeres</i> Skuja	-	-	-	+
<i>Calothrix marchica</i> Lemmermann	-	-	+	-
* <i>Calothrix fusca</i> (Kützing) Bornet & Flahault	-	-	+	-

Trichomes straight, laying parallel and entwined, end cell is conical; cells short, barrel shaped, 4µm broad, 3-4µm long; heterocysts ovate, 5µm broad and 6µm long; akinete developed away from heterocysts, scattered, single or binary, 6-7µm broad and 10-15µm long.

Distribution in studied site: Kalat (36°59'N 59°43' E).

**2. *A. fuelebornii*** Schmidle 1932 (fig. 2B).

Trichomes straight, blue-green, without distinct mucilaginous envelopes. Vegetative cells quadrate, 4µm broad, 5-6µm long; apical cell rounded at the end. Heterocysts barrel shaped, 5µm broad, 8µm long. Akinetes present in one side of the heterocyst, solitary, ellipsoidal, swollen on both sides, about 8µm broad and 27µm long.

Distribution in studied site: Sarrood (36°47'N 59°52' E).

**3. *Nostoc piscinale*** Kützing ex Born. et Flahault 1888

(fig. 2D).

Thallus light blue-green, filaments flexuous, loosely entangled, trichome 3-4µm broad. Vegetative cells subspherical, 3-3.5µm broad and 4 µm long. Heterocysts subspherical or spherical, 4.5-6 µm broad. Akinetes globose, 5-7µm broad, in long chains.

Distribution in studied site: Sarrood (36° 47'N 59° 52' E).

**4. *Nostoc pruniforme*** Ag. ex Bornet et Flahault 1888 (fig. 2F).

Thallus globose or ellipsoid, olivaceous; filaments loosely entangled, 5 µm broad; sheath distinct, hyaline. Vegetative cells short barrel shape, 5µm broad and 2.5-5µm long. Heterocysts spherical, 6-7 µm diameter. Akinetes spherical, 10 µm diameter.

Distribution in studied site: Sarrood (36° 47'N 59° 52' E).




Fig. 2. New records for algal flora of Iran. A, *Anabaena inaequalis*; B, *Anabaena fuellborni*; C, *Nostoc pruniforme*; D, *Nostoc picinale*; E, *Nostoc entophytum*; F, *Calothrix fusca*. (Het: heterocyst; Aka: akinete; spo: spore)




Fig. 3. Heterocystous cyanobacteria observed under optic microscope. A, *Calothrix dolichomeres*; B, *Calothrix elenkinii*; C, *Calothrix marchica*; D, *Anabaena orientalis*; E, *Anabaena torulosa*; F, *Anabaena iyengarii*; G, *Anabaena oscillarioide*; H, *Anabaena vaginicola*; K, *Cyindrospermum sphaericum*; P, Q, R, *Cyindrospermum muscicola*; M, *Nostoc commune*; N, R, *Anabaena portoricensis*.




Fig. 4. Heterocystous cyanobacteria observed under optic microscope. A, B, *Nostoc alatosporum*; C, *Nostoc elliposporum* var. *violaceum*; D, *Nostoc amplissimum*; E, *Nostoc sphaericum*; F, *Nostoc calcicola*; G, *Nostoc verrucosum*; H, *Nostoc punctiforme*; K, *Nostoc linckia*; P, *Nostoc spongiaeforme*; Q, *Nostoc muscorum*; M, N, R, *Nostoc rivulare*.

Table 2. Total percent abundance of heterocystous cyanobacterial genera (Summed up over all location).

Genus	Total no. of species	Percent Abundance
<i>Nostoc</i>	15	50
<i>Anabaena</i>	8	26.66
<i>Cylindrospermum</i>	3	10
<i>Calothrix</i>	4	13.33
<b>Total</b>	<b>30</b>	<b>100</b>

**5. *Nostoc entophytum*** Bornet et Flahault 1888 (fig. 2G).

Thallus small, blue-green, filaments densely entangled; trichomes 2.5-3  $\mu\text{m}$  broad. Vegetative cells short, barrel shap, 2.5  $\mu\text{m}$  broad and 3  $\mu\text{m}$  long. Heterocysts broader than the vegetative cells, 4  $\mu\text{m}$  broad and 3  $\mu\text{m}$  long. Akinetes spherical, 5-6 $\mu\text{m}$  broad.

Distribution in studied site: Amirabad (36° 33'N 60° 5' E).

Rivulariaceae

**6. *Calothrix fusca*** (Kützing) Bornet & Flahault 1886 (fig 2H).

Filaments single, sheath broad, colorless; trichomes 7-8 $\mu\text{m}$  broad. Vegetative cells often shorter than broad, 6 $\mu\text{m}$  broad and 3-7 $\mu\text{m}$  long. Heterocyst basal, hemispherical, single, smaller than the basal cell of the trichome, 4 $\mu\text{m}$  broad.

Distribution in studied sites: Sarood (36° 47'N 59° 52' E).

## DISCUSSION

Heterocystous cyanobacteria have been already studied for their diversity in rice fields of Iran (Shokravi & al. 2002; Siahbalaei & al. 2011), but limited works have been reported on cyanobacteria from paddy fields of northeastern part of this country. In the present study, occurrence of heterocystous cyanobacteria in four study sites of Kalat Naderi district, from Razavi Khorasan Province, was evaluated. The result obtained from this research represent that rice fields of Kalat Naderi contain a rich flora of heterocystous cyanobacteria. Among the identified taxa, members of Nostocacean family were dominant. Similar results were obtained by several researchers and the predominance of Nostocaceae family in rice fields of Iran were reported (Siahbalaei et al. 2011). In our study, *Nostoc* and *Anabaena* from this family had the highest diversity in the soil samples of all sites. Similar to the present study, previous researches reported *Nostoc* as a dominant and most diverse genera of this family in rice fields of Iran (Siahbalaei & al. 2011; Shariatmadari & al. 2013).

Among studied sites, Sarood (Fig. 1, Station 3) with 24 species and Galeno (fig. 1, Station 2) with four species possess the most and the least species variation

respectively. Due to the necessity of thorough soil analysis for conclusions about the species diversity and its reasons, the authors do not have permission to comment definitively about the reason of this diversity in several sites. Nevertheless, in respect to the same weather condition and a slight difference in latitude and longitude of all sites, difference in species composition may be the result of difference in physico-chemical properties of soil in several sites. In the other studies, physical characters such as pH and Electrical conductivity are proposed as the most important factors determining soil floristic composition (Nayak and Prasanna 2007). But the results of some researches indicated that in some of cases, physicochemical factors such as soil pH, EC, total nitrogen, phosphorus and organic carbon, have not significant correlation with the species diversity of cyanobacteria in paddy soils (Shariatmadari & al. 2013). Due to different opinions about this issue, and according to the insufficient data around our studied sites, certain comments require further analysis and ecological studies.

## ACKNOWLEDGMENTS

This study was supported by University of Shahid Beheshti. The authors would like to express their thankfulness to all colleagues in laboratory of phycology for their help.

## REFERENCES

- Anagnostidis, K. & Komárek, J. 1985: Modern approach to the classification system of cyanophytae 1- Introduction.- Arch. Hydrobiol./algolog. Stud. 38-39:291-302.
- Castenholz, R. W. 1989: Subsection IV. Order Nostocales. In Staley, J. T., M. P. Bryant, N. Pefennig & J. G. Holt (eds.). -Bergey's Manual of Systematic Bacteriology 3: 1780-1793.
- Desikachary, T.V. 1959: Cyanophyta, 686 pp.- Indian Council of agricultural Research, New Delhi.
- Ezhilarasi, A. & Anand, N. 2009: Phylogenetic analysis of *Anabaena* spp. (cyanobacteria) using sequences of 16SrRNA gen. -Australian Journal of Basic and Applied Sciences 3(4): 4026-4031.
- Galhano, V., De Figueiredo, D.R., Alves, A., Correia,

- A., Pereira, M. J., Gomes-Laranjo, J. & Peixoto, F. 2011: Morphological, biochemical and molecular characterization of *Anabaena*, *Aphanizomenon* and *Nostoc* strains (Cyanobacteria, Nostocales) isolated from Portuguese freshwater habitats. – *Hydrobiologia* 663: 187-203.
- Komárek, J. & Anagnostidis, K. 1989: Modern approach to the classification system of cyanophytes, 4-Nostocales. – *Arch. Hydrobiol. Suppl. 82/Algological Studies* 56: 247-345.
- Nowruzi, B. & Ahmadimoghadam, A. 2006: Two new Records of heterocystus cyanobacteria (Nostocaceae) from paddy fields of Golestan Province. – *Iran. Journ. Bot.* 11(2): 170-173.
- Prescott, G.W. 1970: *Algae of the western great lake area*. W.M. C. Brown company publishers, 977pp
- Saadatnia, H. & Riahi, H. 2009: Cyanobacteria from paddy fields in Iran as a biofertilizer in rice plants. – *Plant Soil Environ.* 55(5): 207-212.
- Shariatmadari, Z., Riahi, H., Seyed Hashtroudi, M., Ghassempour, A.R. & Aghashariatmadary, Z. 2013: Plant growth promoting cyanobacteria and their distribution in terrestrial habitats of Iran.- *Soil Science and Plant Nutrition* 59: 535-547.
- Shariatmadari, Z & Riahi, H. 2013: A taxonomic study on soil taxa of apoheterocytic cyanoprokaryota from Nostocaceae family in Iran. – *Iran. J. Bot.* 19 (2): 143-152.
- Shokravi, S. 2002: Applied research management of cyanobacteria in Iran: problems and solutions. The first Iranian Congress on Applied Biology, Islamic Azad University, Mashhad, Iran.
- Siahbalaee, R., Afsharzade, S. & Shokravi, S. 2011: New Records of Nostoclean Cyanobacteria from rice fields in the Golestan Province In North-East of Iran. – *Biological Sciences.* 1(2): 50-55.
- Song, T., Martensson, L., Eriksson, T., Zheng, W. & Rasmussen, U. 2005: Biodiversity and seasonal variation of the cyanobacterial assemblage in a rice paddy field in Fujian, China. – *The Federation of European Microbiological Societies Microbiology Ecology* 54:131-140.
- Stanier, R.Y., Kunisawa, R., Mandel, M. & Cohen-Bazire, G. 1971: Purification and properties of unicellular blue- green algae (order Chroococcales). – *Bact. Rev.* 35: 171-205.