

CHROMOSOME COUNT REPORTS OF TWO RARE ENDEMIC SPECIES OF TANACETUM IN IRAN

S. Sadeghian, A. Hatami, E. Jafari & B. Hamzeh'ee

Received 2018. 10. 24; accepted for publication 2019. 04. 18

Sadeghian, S., Hatami, A., Jafari, E. & Hamzeh'ee, B. 2019. 06. 30: Chromosome count reports of two rare endemic species of *Tanacetum* in Iran. -*Iran. J. Bot.* 25 (1): 44-48. Tehran.

The present study is reported somatic chromosome number of two *Tanacetum* species including *T. stapfianum* (Rech.f.) podlech (2 population) and *T. lingulatum* (Boiss.) Bornm. (1 population) for the first time. They belong to *Asteraceae* family and are endemic species from Iran. Two species of *Tanacetum* were collected from their habitats to investigate their karyotypes. Both species were diploid and chromosome numbers were $2n=2x=18$. Ideograms are depicted for each population.

Sara Sadeghian (correspondence <s.sadeghian@areeo.ac.ir>), Ahmad Hatami & Efat Jafari, Research Division of Natural Resources Department, Fars Agricultural and Natural Resources Research and Education Center, AREEO, Shiraz, Iran. -Behnam Hamzehee, Research institute of Forests and Rangelands, Agricultural Research, Education Organization (AREEO), Tehran, Iran.

Key words: *Asteraceae*; Chromosome number; new counting; *Tanacetum*; Iran

گزارش شمارش کروموزومی دو گونه انحصاری نادر از جنس *Tanacetum* در ایران

سارا صادقیان: کارشناس پژوهشی بخش تحقیقات منابع طبیعی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی استان فارس، سازمان تحقیقات، آموزش و ترویج کشاورزی، شیراز، ایران

احمد حاتمی: کارشناس ارشد بخش تحقیقات منابع طبیعی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی استان فارس، سازمان تحقیقات، آموزش و ترویج کشاورزی، شیراز، ایران

عفت جعفری: استادیار پژوهشی بخش تحقیقات منابع طبیعی، مرکز تحقیقات و آموزش کشاورزی و منابع طبیعی استان فارس، سازمان تحقیقات، آموزش و ترویج کشاورزی، شیراز، ایران

بهنام حمزه: دانشیار پژوهشی، مؤسسه تحقیقات جنگلها و مراتع کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران
تحقیق حاضر عدد کروموزومی دو گونه *Tanacetum* به نامهای *T. stapfianum* (Rech. f.) podlech (۲ جمعیت) و *T. lingulatum* (Boiss.) Bornm. (یک جمعیت) را برای اولین بار گزارش می‌کند. این گونه‌ها متعلق به خانواده *Asteraceae* و جزء گیاهان انحصاری ایران می‌باشند. پارامترهای کروموزومی آنها بررسی شد. هر دو گونه دیپلوئید و دارای فرمول کاریوتایی $2n=2x=18$ بودند. ایدیوگرام آنها نیز رسم گردید.

INTRODUCTION

The genus *Tanacetum* L. belongs to *Asteraceae* family. It includes about 160 species in the world and is the third largest genus of tribe *Anthemideae*, distributed in Asia, North Africa, North America and

Europe (Oberprieler & al. 2007). This genus has 54 species in the area of Flora Iranica (podlech 1986) and has 31 species in Iran of which 16 species are endemics (Mozaffarian 2005, 2008). Many karyological data concerning chromosome numbers have already been

published in *Tanacetum* and the most common basic chromosome number in the genus was found to be $x=9$. There are some reports of ploidy levels in some species. Javadi (2017) studied 14 populations of *Tanacetum polycephalum* and *Tanacetum parthenium* from different parts of Iran and reported that in all of them the basic chromosome number was $x=9$ and showed two ploidy levels (diploid and tetraploid). Sonboli & al. (2011) studied *T. fisherae* from Kerman province and showed chromosome number $2n=44+1B$ that indicates a pentaploid level ($5x$) based on $x=9$, which is a new ploidy level for this genus. Chehregani & al. (2011) recorded various levels of polyploidy within 14 populations of *T. polycephalum*. The number of chromosome was 18 (diploid), 36 (tetraploid) and 54 (hexaploid). Chehregani & al. (2012) also reported evolutionary karyotypic variation in *Tanacetum parthenium* populations in Iran and confirmed the existence of two levels of ploidy, diploid ($2n=2x=18$) and tetraploid ($2n=4x=36$) in the species. The chromosome numbers for the endemic species, *Tanacetum stapfianum* and *Tanacetum lingulatum* have not been reported yet. In this study we investigated the karyotypes of these two species for the first time.

MATERIALS AND METHODS

Two species *Tanacetum stapfianum* (two populations) and *Tanacetum lingulatum* (one population) have been studied (table 1). Vouchers are deposited in the Herbarium of Fars Agricultural and

Natural Resources Research and Education Centre of Iran.

For cytological study, rootlets were collected from germinated seeds on wet filter paper in petri dishes at 22°C temperature, when they reached 1–1.5 cm in length, rootlets were separated. The root tips meristems treated with 0.5% saturated of α -Bromo naphthalene at 4°C for 4-5 h. Then were fixed in 10% formaldehyde and chromium trioxide (1: 1) for 20 to 24 h at 4°C. Then the root tips were rinsed for 1 h in distilled water. Hydrolysis was carried out with NaOH (1 normal) at 60°C for 20-25 minutes and used hematoxylin-iron for chromosome staining for 1-2 h at room temperature. Root tips were squashed in a droplet of 45% acetic acid. The preparations were observed with an optical microscope (BH2 Olympus supplemented Digital color video camera) at a magnification of 2000x.

RESULTS

There was no difference between basic chromosome number of the two species ($x=9$). The somatic chromosome numbers ($2n$), karyotype formula and parameters for the studied species and populations are summarized in table 2. Two species were diploid with $2n=2x=18$. The chromosomes were mostly metacentric (m) (Levan & al. 1964), (fig. 1) and two populations of *T. stapfianum* are categorized in type 1A and *T. lingulatum* species is categorized in type 2A (table 1). Ideograms are depicted for each species and population (fig. 2).

Table 1. Karyotype characters of two *Tanacetum* species. $2n$: Diploid chromosome numbers; A_1 : intrachromosome asymmetry index; A_2 : interchromosome asymmetry index; TF%: total form percentage; DRL: difference of relative length; VRC: value of relative chromatin; SC: symmetry classes of Stebbins and K.F.: karyotype formula.

species	locality	$2n$	A_1	A_2	VRC	DRL	%TF	SC	K.F.
<i>Tanacetum stapfianum</i> (Dasht-e Arjan)	Iran, Fars province, west of Shiraz, Dasht-e Arjan, 12812.	$2x=18$	0.19	0.13	8.20	4.66	44.75	1A	$7m+2sm$
<i>T. stapfianum</i> (Bamoo protected area)	Iran, Fars province, north of Shiraz, Bamoo protected region, 12813.	$2x=18$	0.21	0.13	8.90	4.51	43.91	1A	9m
<i>T. lingulatum</i>	Iran, Fars province, east of Shiraz, Sarvestan, 12814.	$2n=18$	0.26	0.17	6.09	5.55	41.88	2A	$8m+1sm$

Table 2. Mean of chromosomes analysis of two *Tanacetum* species. TL: total length of chromosome; LA: long arm; SA: short arm; AR: arm ratio; CI: centromeric index; DRL: difference of relative length; TF%: total form percentage; %SA: Relative length percentage of short arm; %LA: Relative length percentage of long arm.

species	TL	LA	SA	AR	CI	DRL	%TF	%SA	%LA
<i>Tanacetum stapfianum</i> (Dasht-e Arjan)	8.20	4.53	3.67	1.27	0.45	4.66	44.75	4.97	6.13
<i>T. stapfianum</i> (Bamoo protected area)	8.87	4.96	3.91	1.28	0.44	4.51	43.91	4.88	6.19
<i>T. lingulatum</i>	6.09	3.54	2.55	1.45	0.42	5.55	41.88	4.65	6.04




Fig. 1. Somatic metaphases in *Tanacetum* species. a, *T. stapfianum* (population of Dasht-e Arjan); b, *T. stapfianum* (population of Bamoo protected area) and c, *T. lingulatum*.




Fig. 2. Ideograms of *Tanacetum* species. a, *T. stapfianum* (population of Dasht-e Arjan); b, *T. stapfianum* (population of Bamoo protected area) and c: *T. lingulatum*.

REFERENCES

- Chehregani A., Mohsenzadeh F., Moazen F. 2012: Variation of chromosome numbers in *Tanacetum parthenium* Schultz Bip. (Asteraceae) in Iran. - *Chromosome Botany*. 7: 97-100.
- Chehregani A., Atri M., Sarmadi J. Asgari M. 2011: Chromosome number variation in *Tanacetum polycephalum* Schultz bip. (L.) (Asteraceae) in west of Iran. - *Caryologia* 64(3): 303-308.
- Javadi H. 2017: Karyological data of *Tanacetum polycephalum* Schultz-bip. And *T. parthenium* Schultz-bip. (Asteraceae) populations. - *Journal of Cell and Molecular Research*. 9 (2): 59-66.
- Levan A. K., Fredga K., Sandberg A. A. 1964: Nomenclature for centromic position on chromosomes. - *Hereditas*. 52: 201-220.
- Mozaffarian V. 2005: Notes on the tribe *Anthemideae* (compositae), new species, new records and new combinations for Iran. - *Iran J. Bot.* 11 (1): 115-127.
- Mozaffarian V. 2008: Compositae, *Anthemideae* and *Echinopeae* in M. Assadi & al. (eds.), *Flora of Iran* 59: 134-198. - Research Institute of Forests and Rangelands, Tehran.
- Oberprieler C, Vogt R, Watson LE. 2007: *Anthemideae*. In *The families and genera of vascular plants VIII*. Vol 8, Berlin, p. 342-374.
- Podlech D. 1986: Compositae VII: *Tanacetum* in: Rechinger KH (ed.), *Flora Iranica*: vol. 158. Akademische Druck- und Verlagsanstalt, Graz.
- Sonboli A., Olanj N., Pourmirzaei A. 2011: Biosystematics and phylogeny of *Tanacetum fisherae*, a new record from Iran. - *Rostaniha*, 12(2): 165-175.
- Stebbins G. L. 1971: *Chromosomal evolution in higher plants*. Edward Arnold Publisher, London, Ltd. Pp. 216.