

KARYOTYPE ANALYSIS OF THREE SPECIES FROM THE WESTERN REGIONS OF IRAN

A. Shariat, A. Jalili, S. S. Mirzadeh Vaghefi & M. Khanhasani

Received 2021. 12. 26; accepted for publication 2022. 06. 07

Shariat, A., Jalili, A., Mirzadeh Vaghefi, S. S. & Khanhasani, M. 2022. 06. 30: Karyotype analysis of three species from the western regions of Iran. *Iran. J. Bot.* 28 (1): 64-68. Tehran.

Three plant species belonging to three genera of Apiaceae, Boraginaceae & Poaceae were collected from wetlands in western regions of Iran. The chromosomal count of *Leiotulus porphyrodiscus* (Stapf & Wettst.) Pimenov & Ostr. ($2n = 22$) is reported for the first time, and the chromosome count of *Arrhenatherum elatius* (L.) P. Beauv. ex J. Presl & C. Presl ($2n = 28$) is reported for the first time for the flora of Iran. The number of chromosomes in *Echium italicum* L. ($2n = 16$) is reported from Kermanshah Province. Karyomorphological parameters such as karyotypic formula, Romero-Zarco indices, Stebbins class, and other karyotypic symmetry indices were determined for all species.

Anahita Shariat (correspondence <shariat@rifr-ac.ir>), Adel Jalili, and Saeedeh Sadat Mirzadeh Vaghefi, Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO). Tehran, I.R. Iran. -Masoumeh Khanhasani, Kermanshah Agricultural and Natural Resources Research and Education Centre, Agricultural Research, Education and Extension Organization (AREEO), Kermanshah, I.R. Iran.

Keywords: Chromosome counts; ideogram; symmetry indices; karyotype; flora of Iran

بررسی کاربوتیبی سه گونه از مناطق غربی ایران

آناهیتا شریعت: پژوهشگر مؤسسه تحقیقات جنگلها و مراتع کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران
عادل جلیلی: استاد پژوهش مؤسسه تحقیقات جنگلها و مراتع کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران
سعیده سادات میرزاده واقفی: استادیار پژوهش مؤسسه تحقیقات جنگلها و مراتع کشور، سازمان تحقیقات، آموزش و ترویج کشاورزی، تهران، ایران

معصومه خان حسینی: مربی پژوهش مرکز تحقیقات کشاورزی و منابع طبیعی استان کرمانشاه، سازمان تحقیقات، آموزش و ترویج کشاورزی، کرمانشاه، ایران

سه گونه گیاهی متعلق به سه تیره Apiaceae, Boraginaceae, Poaceae از مناطق ماندابی غرب کشور جمع آوری گردید. شمارش کروموزومی برای گونه *Leiotulus porphyrodiscus* (Stapf & Wettst.) Pimenov & Ostr. ($2n = 22$) برای اولین بار و شمارش کروموزومی گونه *Echium italicum* در گونه *Arrhenatherum elatius* (L.) P. Beauv. ex J. Presl & C. Presl ($2n = 28$) برای اولین بار در فلور ایران و در گونه *L. nitens* ($2n = 16$) از استان کرمانشاه گزارش می گردند. پارامترهای کاربومورفولوژیکی از جمله فرمول کاربوتیبی، شاخص های روموزرکو، کلاس Stebbins و سایر شاخص های تقارن کاربوتیبی برای همه گونه ها تعیین شد.

INTRODUCTION

Historically, the study of wetland habitats in Iran has primarily focused on Anzali, Gavkhooni, and

Jazmourian wetlands while the scattered mountain wetland conditions have been less studied (Khanhasani & al. 2021). Plants that grow in such ecosystems are

called hydrophytic plants. These plants have adapted to special ecological conditions, such as inorganic soils or soft soils that are flooded with poor drainage. Aquatic plants can complete their reproductive cycle in water. However, in addition to the mentioned aquatic plants, other aquatic plants that need a dry environment to complete their life cycle are also considered real aquatic plants. (Den Hartog & van der Velde 1988). The Kermanshah Province occupies about 1.5 percent of Iran with an area of 25000 square kilometers. In the Zagros Mountain range with mountains such as Shahu, Dalakhani, Sefidkooh, differences in altitudes, and climates, have led to a significant diversity of plant species. There are more than one hundred open springs (mirage) as the source of rivers or their tributaries and five permanent rivers and several wetlands and reservoirs in Kermanshah province. Khanhasani & al. (2021) with the aim of preparing a distribution map of wetland plants in Kermanshah province, studied 31 wetland habitats and reported a total of 288 plant species. Few studies have been done on the cytogenetic characteristics of wetland plants in Iran. Shariat & al. (2021a) reported the karyotypic characteristics of 12 wetland species in Kermanshah province. In other research, Mirzadeh Vaghefi and Jalili (2019) represented a chromosome count of 14 species from six families of Iranian wetlands. The present study is part of a comprehensive project to determine the

chromosome number of plants in wetland habitats of Iran. It is worth noting that some studied species were not wetlands, but due to favorable environmental conditions around wetlands, they have grown there. A major goal of this study is to strengthen the information on Iran's plant cytogenetic database.

MATERIALS AND METHODS

Plants were collected from two wetland areas of Kermanshah province named Hashilan, and Gamasiab. Plant samples were identified and herbarium specimens were prepared for examined species. For karyotypic studies, the seeds were first germinated in sterilized Petri dishes. The roots were pretreated with α -Bromonaphthalene solution (2 h), then fixed in Carnoy solution for 24 hours (3 ethanol: 1 glacial acetic acid v/v). The roots were kept in 70% alcohol until the time of testing. The roots were then hydrolyzed using 1 N hydrochloric acids at 60 °C for 10 minutes and stained with 4% hematoxylin for 6 hours at 60 °C and finally squashed in 45% (v/v) acetic acid (Shariat & al. 2013; Shariat and Sefidkon 2021b). An optical microscope (Nikon Coolpix P90 digital camera interfaced to a BH2- RFCA Olympus microscope) was used for mitotic studies. Karyotypic parameters and asymmetry indices were measured by Ideokar software (Mirzaghaderi & Marzangi 2015) as described in table 1.

Table 1. Measures and formulae of karyotypic parameters and asymmetry indices. Abbreviations: L, length of long arm; S, length of short arm; I, homologous group number; s, standard deviation; X, mean; n, haploid chromosome number of an individual or taxon.

No	Karyotypic parameters	Formula	Reference
1	Total chromosome length of the haploid complement	$HCL = \sum CL$	Levan & al. (1964)
2	Total form percentage	$TF\% = (\sum S / \sum CL) * 100$	Huziwarra (1962)
3	Arano index of karyotype asymmetry	$AsK\% = (\sum L / \sum CL) * 100$	Arano (1963)
4	Intrachromosomal asymmetry index	$A1 = \sum_{i=1}^n (S_i / L_i) / n$	Romero-Zarco (1986)
5	Interchromosomal asymmetry index	$A2 = S_{CL} / X_{CL}$	Romero-Zarco (1986),
6	Symmetry index	$S\% = (CL_{min} / CL_{max}) * 100$	Watanabe & al. (1999)
7	Mean centromeric index	$X_{Cl} = \sum CL / n$	Paszko (2006)
8	Degree of karyotype asymmetry	$A = \sum_{i=1}^n [(L_i - S_i) / (L_i + S_i)] / n$	Watanabe & al. (1999)
9	Mean centromeric asymmetry	$X_{CA} = A * 100$	Paszko (2006)
10	Coefficient of variation of chromosome length	$CV_{CL} = (S_{CL} / X_{CL}) * 100 = A_2 * 100$	Paszko (2006)
11	Coefficient of variation of centromeric index	$CV_{Cl} = (S_{Cl} / X_{Cl}) * 100$	Paszko (2006)
12	Asymmetry index	$AI = (CV_{CL} * CV_{Cl}) / 100$	Paszko, (2006)

RESULTS AND DISCUSSION

Apiaceae

Leiotulus porphyrodiscus (Stapf & Wettst.) Pimenov & Ostr. ($2n = 22$), (Fig. 1, Table 2)

Iran: Kermanshah Province, Hashilan, 1316 m, 03.08.2021, 34°35'4.9"N, 46°53'46"E. Khanhasani and Rahimi 8899 (RANK).

The karyotypic formula represented 13 median, seven metacentric, and two subterminal chromosomes (13 m + 7 sm + 2 st). The karyotype is relatively symmetrical and classified as Stebbins 1B symmetry. The total chromosome length of the haploid complement is 69.8 μm (Table 2). This is the first chromosome number report of this species.

The chromosome counts from the other species of the genus, including *L. pastinacifolius* (Boiss. & Balansa) M.G. Pimenov & T.A. Ostroumova, *L. secacul* (Mill.) M.G. Pimenov & T.A. Ostroumova, and *L. dasyanthus* (K. Koch) M.G. Pimenov & T.A. Ostroumova from the Mediterranean region have previously been reported as $2n = 20$, $2n = 44$, and $2n = 22$ respectively (Shner 2004 & 2005).

Boraginaceae

Echium italicum L. ($2n = 16$), (Fig. 1, Table 2)

Iran: Kermanshah Province, Hashilan, 1316 m, 03.08.2021, 34°35'4.9"N, 46°53'46"E. Khanhasani and Rahimi 8234 (RANK).

This species has a symmetrical karyotype and fell in 1B of Stebbins classification. The karyotype formula is 2M + 10m + 2sm + 2st. The total chromosome length of the haploid complement is 37.5 μm . A value of TF = 44.1% indicates intrachromosomal differences. A higher value of TF indicates greater symmetry of

chromosomes, while S = 32.7% indicates the difference between the minimum and maximum length of chromosomes. In addition to the two mentioned indices, Romero-Zarco indices are in agreement with the values of S% and TF%, or in other words, with the increase of TF, the value of A1 decreases and with the decrease of S, the value of A2 increases. It can be concluded that in *Echium italicum*, the difference between the short and long arms is small, while there is a significant difference between the length of the longest and shortest chromosomes, or in other words, the intrachromosomal differences are minimal, while the interchromosomal differences are significant. The results of this study are in agreement with the results of Azoush & al. 2014 and Ghaffari 1996, who worked on the populations of Tehran (Fasham) and Karaj, respectively.

Poaceae

Arrhenatherum elatius (L.) P. Beauv. ex J. Presl & C. Presl ($2n = 28$), (Fig. 1, Table 2)

Iran: Kermanshah Province, Gamasiab, 1290 m, 14.07.2021, 34°27'8.3"N, 47°26'19.8"E. Khanhasani and Rahimi 8122 (RANK).

The karyotype formula is 3M + 17m + 6sm + 2st. This species has a symmetrical karyotype and fell in 1B of Stebbins classification. The total chromosome length of the haploid complement is 140.4 μm (Table 1). This is the first report on the chromosome number of *Arrhenatherum elatius* for the flora of Iran. Our result is consistent with the results of previous counts from Norway (Engelskjön 1979) and Armenia (Ghukasyan 2004).

Table 2. Chromosomal parameters and karyotype asymmetry indices of 12 species from Kermanshah Province.

Species	HCL	TF	AsK%	S%	Xci	A	XcA	CVcl
<i>Leiotulus porphyrodiscus</i>	69.8	39.4	60.6	41.5	0.38	0.23	23.3	17.5
<i>Echium italicum</i>	37.5	44.1	55.9	32.7	0.44	0.12	11.6	36.2
<i>Arrhenatherum elatius</i>	140.4	40.5	59.5	30.8	0.41	0.18	18.2	26.8

Table 2. Continued.

Species	CVci	AI	Stebbins Class	Asymmetry indices (Romero-Zarco)		
				A1	A2	KF
<i>Leiotulus porphyrodiscus</i>	21.7	80.8	1B	0.35	0.18	13 m + 7 sm + 2 st
<i>Echium italicum</i>	16.4	221.0	1B	0.18	0.36	2M + 10m + 2sm + 2st
<i>Arrhenatherum elatius</i>	18.8	142.1	1B	0.28	0.27	3M + 17m + 6sm + 2st




Fig. 1. Karyotypes and ideograms of the examined species. a, *Leiotulus porphyrodiscus* ($2n = 22$); b, *Echium italicum* ($2n = 16$); c, *Arrhenathrum elatius* ($2n = 28$); Bar = 10 μ m.

ACKNOWLEDGEMENTS

This research was supported by the Research Institute of Forests and Rangelands (RIFR).

REFERENCES

- Arano, H. 1963: Cytological studies in subfamily *Carduoideae* (Compositae) of Japan XI.-Bot. Mag. 76: 32-39.
- Azoush, S., Kazemitabar, S. K. & Heidari, P. 2014: Polyploidy induction in Iranian Borage (*Echium amoenum* L.) by colchicine treatment. -Biharian Biologist. 8 (2): 87-89.
- Den Hartog, C. & van der Velde, G. 1988: Structural aspects of aquatic plant communities. In: Handbook of vegetation science, Symones, J.J. (eds.), vol. 5: Vegetation of inland water. Dr. W. Junk Publishers, Dordrecht: 113-153.
- Engelskjøn, T. 1979: Chromosome numbers in vascular plants from Norway, including Svalbard. -Opera Bot. 52: 1-38.
- Ghaffari, S. M. 1996: Chromosome studies in some species of Boraginaceae from Iran. -Iran. J. Bot. 7 (1): 81-93.
- Ghukasyan. 2004: Extent of karyological study of Armenian grasses (Poaceae). Fl. Rastitel'n. Rastitel'n. Resursy Armenii 15: 85-89.
- Huziwaru, Y. 1962: Karyotype analysis in some genera of Compositae. VIII. Further studies on the

- chromosomes of Aster. -Am J of Bot. 49 (2):116-119.
- Khanhasani, M., Jalili, A., Khodakarami, Y. & Jalilian, N. 2021: Wetland flora of Kermanshah Province, - Iran. Nova Biologica Reperta. 8 (2): 154-172. (In Persian).
- Levan, A., Fredga, K. & Sandberg A. A. 1964: Nomenclature for centromeric position on chromosomes. -Hereditas. 52 (2): 201-220.
- MirzadehVaghefi, S., Jalili, A. 2019: Chromosome numbers of some vascular plant species from Iran. - Iran. J. Bot. 25 (2): 140-144.
- Mirzaghaderi, G., & Marzangi, K. 2015: IdeoKar: an ideogram constructing and karyotype analyzing software. -Caryol. 68 (1), 31-35. <https://doi.org/10.1080/00087114.2014.998526>
- Paszko, B. 2006: A critical review and a new proposal of karyotype asymmetry indices. -Plant Syst Evol. 258 (1): 39-48.
- Romero-Zarco, C. 1986: A new method for estimating karyotype asymmetry. -Taxon. 35 (3): 526-530.
- Shariat, A., Jalili, A., Mirzadeh Vaghefi, S. S. & Khan Hasani, M. 2021a: Chromosome counts of twelve vascular plant species from Kermanshah province. Iran J Bot. 27 (2): 1-12.
- Shariat, A. & Sefidkon, F. 2021b: Tetraploid induction in savory (*Satureja khuzistanica*): cytological, morphological, phytochemical and physiological changes. -Plant Cell, Tissue, and Organ Cult. 1-13.
- Shariat, A., Karimzadeh, G. & Assareh, M. H. 2013: Karyology of Iranian Endemic *Satureja* (Lamiaceae) Species. -Cytol. 78 (3): 305-312.
- Shner, J. V. 2004: Mediterranean chromosome number reports 14 (1391-1409). -Fl. Medit. 14: 435-447.
- Shner, J. V. 2005: Mediterranean chromosome number reports 15 (1438-1442). -Fl. Medit. 15: 710-716.
- Watanabe, K., Yahara, T., Denda, T. & Kosuge, K. 1999: Chromosomal evolution in the genus *Brachyscome* (Asteraceae, Astereae): statistical tests regarding the correlation between changes in karyotype and habit using phylogenetic information. -J Plant Res. 112 (2): 145-161.